

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/317393655>

Extracorporeal shockwave: mechanisms of action and physiological aspects for cellulite, body shaping, and localized fat—Systematic review

Article in *Journal of Cosmetic and Laser Therapy* · June 2017

DOI: 10.1080/14764172.2017.1334928

CITATIONS

14

READS

1,350

8 authors, including:



Débora Aparecida Oliveira Modena
University of Campinas

28 PUBLICATIONS 46 CITATIONS

[SEE PROFILE](#)



Renata Michelini Guidi
University of Campinas

14 PUBLICATIONS 59 CITATIONS

[SEE PROFILE](#)



Estela Sant'Ana
www.cienciaeestetica.com.br

9 PUBLICATIONS 61 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Fisioterapia Respiratória [View project](#)



Obesidade [View project](#)



Extracorporeal shockwave: mechanisms of action and physiological aspects for cellulite, body shaping and localized fat – systematic review.

Journal:	<i>Journal of Cosmetic and Laser Therapy</i>
Manuscript ID	Draft
Manuscript Categories:	Reviews of Treatment Studies
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Modena, Débora; Universidade Estadual de Campinas, Surgery Department; Ibramed Research Institute, Study Group in Technology Applied to Health, Brazil</p> <p>Silva, Caroline; Universidade Estadual Paulista Julio de Mesquita Filho - Campus de Marilia, Human Development and Technologies; Ibramed Research Institute, Study Group in Technology Applied to Health, Brazil</p> <p>Grecco, Clovis; Ibramed Research Institute, Research Development and Innovation</p> <p>Guidi, Renata; Universidade Estadual de Campinas, Biomedical Engineering Department; Ibramed Research Institute, Study Group in Technology Applied to Health, Brazil.</p> <p>Moreira, Renata; Ibramed Research Institute, Study Group in Technology Applied to Health, Brazil</p> <p>Coelho, Andresa; Ibramed Research Institute, Study Group in Technology Applied to Health, Brazil</p> <p>Sant'Ana, Estela; Ibramed Research Institute, Study Group in Technology Applied to Health, Brazil.</p> <p>Souza, José; Ibramed Research Institute, Study Group in Technology Applied to Health, Brazil.</p>
Keyword:	Cellulite, acoustic wave therapy, extracorporeal shockwaves, shockwaves cellulite

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Title Page

*Extracorporeal shockwave: mechanisms of action and physiological aspects for
cellulite, body shaping and localized fat – systematic review.*

Physiology extracorporeal shock waves

MSc. Débora A Oliveira Modena^{a,c}, PT. Caroline Nogueira da Silva^{b,c}, PhD. Clovis Grecco^c, PT. Renata Michelini Guidi^{c,d,e}, PT. Renata Gomes Moreira^c, Andresa A Coelho^c, PhD. Estela Sant'Ana^c, José Ricardo de Souza^{c,e}.

^a*Of Surgery Department, Faculty of Medical Sciences, University of Campinas (Unicamp), Brazil.*

^b*Human Development and Technologies, Universidade Estadual Paulista (UNESP), Brazil.*

^c*Ibramed Research Institute: Study Group in Technology Applied to Health, Brazil.*

^d*Biomedical Engineering Department, Faculty of Electrical Engineering and Computing, University of Campinas (Unicamp), Brazil.*

^e*Centro de Estudos e Formação Avançada Ibramed (CEFAl), Brazil.*

Corresponding Author: MSc. Débora A Oliveira Modena

de.modena@ibramed.com.br

Av. Dr. Carlos Burgos, 2800 - Jardim Itália, Amparo - SP, 13901-080

Contributions of all authors

MSc. Débora A Oliveira Modena^{a,c}, search for articles in the database, articles analysis, manuscript writing.

PT. Caroline Nogueira da Silva^{b,c}, search of articles in the database, articles analysis, manuscript revision, contributions in writing the manuscript.

PhD. Clovis Grecco^c, manuscript revision, contributions in writing the manuscript.

PT. Renata Michelini Guidi^{c,d,e} manuscript revision, contributions in writing the manuscript.

PT. Renata Gomes Moreira^c, manuscript revision, contributions in writing the manuscript.

Andresa A Coelho^c, manuscript revision, contributions in writing the manuscript.

PhD. Estela Sant'Ana^c, manuscript revision, contributions in writing the manuscript.

José Ricardo de Souza^{c,e}. manuscript revision, contributions in writing the manuscript.

1
2
3
4
5 *Extracorporeal shockwave: mechanisms of action and physiological aspects for*
6
7 *cellulite, body shaping and localized fat – systematic review.*
8

9
10 **Abstract:** Extracorporeal Shockwave Therapy – ESWT has had a wide use in
11 rehabilitation, and has presented positive effects in the treatment of unaesthetic
12 affections. The objective of the present study was to search, in the literature, the
13 mechanisms of action and the physiological aspects of shockwaves acting on the
14 biological tissue to improve the condition of cellulite and localized fat. A systematic
15 review of the literature was carried out in the period from September 2016 to February
16 2017 based on the bibliographic databases Lilacs, MedLine, PubMed and Scielo.
17 Fifteen articles were identified in that systematic review, three of which were excluded
18 because they did not make the complete access to the article available or the theme
19 investigated did not encompass the objective of the study. The revision demonstrated
20 that extracorporeal shockwaves present relevant effects on the biological tissue, which
21 leads to the restructuring of skin properties and subcutaneous tissue, clinically
22 improving the aspects of cellulite and localized fat.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

38 **Keywords:** cellulite, acoustic wave therapy, extracorporeal shockwaves, shockwaves
39 cellulite, shockwaves subcutaneous fat.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

Growing technological advances in technology have promoted the development of new electromedical devices that can stimulate new therapeutic modalities. The Extracorporeal Shock Wave Therapy - ESWT is an example of technological evolution that opens the possibility for a non-invasive therapeutic modality, validated for use in the healthcare sector, indicated for the treatment of musculoskeletal disorders and which has recently been explored in the scope of aesthetic affections. Even with the growing use of this technology, its mechanisms of action have not yet been clearly established (1,2).

The use of ESWT had its beginnings in the early 70s in Extracorporeal Lithotripsy, then becoming a golden standard for the treatment of gallstone, renal calculi, ureteral calculi and vesical calculi [3,4]. During Lithotripsy treatment for lower ureteral calculi, physicians observed that some patients presented increase in bone density in the pelvic region. Based on this observation, they began to use equipment designed for Lithotripsy for the treatment of orthopedic dysfunctions, originating a new technology that allowed the grading of depth and intensity with which shockwaves penetrate biological tissues (4,5).

In 1997 in Vienna, Austria, the European Society for Musculoskeletal Shockwave (ESMST) was founded, with the objective of promoting the research and development of therapy in Europe. In Brazil, the use of shockwaves as an orthopedic therapy started in 1998. In 2001, ESWT was approved by the Federal Drug Administration (FDA), for the treatment of chronic plantar fasciitis in the United States (EUA) (5,6).

The possibility of the use of ESWT in aesthetic treatments emerged from the observations of surgical results, when women suffering from muscle disorders, after

1
2
3 receiving hip prosthesis, reported an enhancement in mobility and pain, an improvement
4
5 in the aspect of the skin and body contour, suggesting that there was a decrease in the
6
7 body circumference in the treated area (4,7,8).
8

9
10 The ESWT technology appeared during World War II when submarines
11
12 were attacked by bombs and remained intact, and the sailors on board presented serious
13
14 trauma with visceral and pulmonary lesions without external lesions. These lesions were
15
16 attributed to the shockwaves, which went through the submarine walls and propagated
17
18 amongst the sailors. In everyday life, shockwaves can be perceived when there is
19
20 thunder, which occurs due to the intense heating and the rapid expansion of the air
21
22 surrounding the lightning (4,9).
23

24
25 ESWT is characterized by a high pressure pulse (80 MPa) in a time interval
26
27 of extremely short duration (in the order of nanoseconds), which produces mechanical
28
29 pressure waves that promote cavitation in liquid medium (gas bubbles in fluid medium)
30
31 and increase in local temperature. These shockwaves propagate in the target tissue,
32
33 generating essential mechanisms of action such as the mechanical impact in the tissues
34
35 and formation of microbubbles, which favors the achievement of the therapeutic
36
37 objective (4).
38
39

40
41 For therapeutic purposes, there are four types of shockwave generators: The
42
43 electro-hydraulic system, the piezoelectric system, the electromagnetic system and the
44
45 pneumatic system, the latter two being the most frequently used in rehabilitation and
46
47 aesthetics (6,10). The electromagnetic generator presents two coils inside that generate a
48
49 magnetic field when run through by electricity – the action of the field causes the
50
51 projectile, situated inside the applicator, to move quickly and collide with the tip of the
52
53 applicator. In the collision of the bullet, there is the transfer of energy from the
54
55 projectile to the tip of the applicator, which then transfers the energy received in the
56
57
58
59
60

1
2
3 form of mechanical waves to the treatment area. In the pneumatic generator, differently,
4
5 the projectile is displaced towards the tip by the pressure exerted by the compressed air
6
7 (5,6,10) (Figure 1).
8

9
10 ESWT equipment can be built in such a manner that it emits two types of
11
12 waves, classified as focal waves (more intense and profound), directed at the target
13
14 tissue, or radial waves (less intense and more superficial) diverge over the treatment
15
16 area (5,11). There is equipment which offer the two types of wave and other that use
17
18 only one which is adequate to the therapeutic objective. The ESWT energy can also be
19
20 classified as low, medium and high, and that favors the field of action because each
21
22 energy reaches structures like muscles, tendon and bones in different depths and forms
23
24 (7,8,12).
25
26

27
28 Considering that this technology is relatively new in the aesthetic area, there
29
30 are few studies that discuss the interaction between ESWT and the biological tissues for
31
32 this purpose, which makes decision-making about the use of this technology difficult
33
34 based on evidence. Therefore, the objective of this study was to present a revision of the
35
36 literature about the technical aspects, mechanisms of action and physiological effects
37
38 that involve the use of extracorporeal shockwave therapy in aesthetic alterations.
39
40
41

42 ***Material and Methods***

43
44
45

46 This is a revision of the literature carried out from September 2016 to February based
47
48 on bibliographic databases Lilacs, MedLine, PubMed and Scielo. As the theme is
49
50 relatively recent, there is difference in the indexing processes in the bibliographic
51
52 databases; therefore, the search for free themes, without the use of the controlled
53
54 vocabulary (descriptors) was the option chosen. Using this strategy, there was an upturn
55
56 of a greater number of references, guaranteeing the detection of most of the published
57
58
59
60

1
2
3 studies within pre-established criteria. Terms such as cellulite, acoustic wave therapy,
4
5 extracorporeal shockwaves, shockwaves cellulite, shockwaves subcutaneous fat were
6
7 combined. The selected studies included the extracorporeal shockwave theme and the
8
9 discussion of the mechanisms of action and physiological effects in the treatment of
10
11 aesthetic alterations.
12

13
14 All of the original articles indexed about the action of ESWT on cellulite
15
16 and localized fat written in English published between 2005 and 2017 with
17
18 experimental (clinical trials, randomized or not) or observational (case studies, case-
19
20 control, cross-sectional studies and before-and-after treatment studies) approaches
21
22 carried out in humans were included. Experimental studies were excluded.
23

24
25 After checking titles, authors and abstracts, with the objective of avoiding
26
27 the repetition of publications, the availability of complete access and the theme were
28
29 investigated, and an evaluation of articles by peers, specialists in the investigated area,
30
31 was carried out.
32

33
34 After the inclusion of the articles, an analytical reading of each study was
35
36 performed, as well as a record of concept, method used for evaluation, classification of
37
38 ESWT use in the pre-established area, in the mechanisms of action, the physiological
39
40 effects and last but not least, a comparative analysis between the findings and the
41
42 theoretical discussion.
43
44

45 46 **Results**

47
48 In the systematic review, 15 papers were identified, three of them were excluded
49
50 because not providing complete access availability, or they were animal studies or the
51
52 theme investigated did not encompass the objective of the study. The selected papers
53
54 are presented according to Table 1.
55
56
57
58
59
60

Table 1. Published studies about the action of ESWT on cellulite and localized fat.

ESWT: Extracorporeal Shockwave Therapy

Discussion

The objective of the present study was to search for literary evidence about the physiological effects of ESWT in biological tissues in aesthetic disorders. ESWT has continuously been indicated for the treatment of cellulite, also known as Gynoid Lipodystrophy (12-14). Cellulite is an aesthetic alteration which attacks the thighs, gluteus, abdomen and upper region of the arms. It is a multifactor disorder with increase of adipose tissue that causes cellular dystrophy, dysfunctions of the hydraulic metabolism with blood congestion, tissue fibrosis and proliferation of fibroblasts around the adipose cells (1,12,14). Cellulite is more common in women, because they present about 21 to 22 billion adipose cells, whereas men present about 17 to 18 million. This difference in the quantity of adipocytes causes men to have a smaller ability to store fat in relation to the female sex. Besides, the subcutaneous conjunctive tissue of women presents some particularities such as the fine fibrous septum, directed vertically in the cutaneous surface, causing the adipose cells to present a regular form and the projection of these structures become superficial reaching the dermis, giving the skin the “orange peel” appearance (9,15-17).

Several treatments for cellulite have been developed in the last decades, with the objective of promoting the improvement of blood and lymphatic circulation using physical treatments, pharmacotherapy and treatments in association with electromedical equipment. However, there is not any one treatment that is completely effective (16–20).

1
2
3 The first ESWT study for the treatment of cellulite was carried out in 2005,
4
5 in a clinical study with 26 women with average age of 45 years, which compared the
6
7 independent application of ESWT and in combination with decongestive therapy for
8
9 two weeks. Six sessions of ESWT were performed with 1.000 shots in only one thigh
10
11 and the decongestive therapy was performed in both thighs. The authors concluded that
12
13 one single application of ESWT can significantly decrease the levels of Plasma
14
15 Malondialdehyde (MDA), a biomarker of the level of oxidative stress, which would
16
17 cause improvement of the biomechanical properties of the skin (19). Oxidative stress is
18
19 generated when there is imbalance in the production and tamponade of free radicals,
20
21 biochemical physiological alterations during metabolic processes or due to external
22
23 factors such as poor nutrition, exhausting exercise, psychological stress, and tobacco
24
25 use, among others. As we suffer the influence of these factors, the cell walls and their
26
27 molecules are attacked, causing a continuous and aggressive imbalance, which
28
29 unleashes a cascade of events that induce the cellular aging. When the oxidative stress
30
31 reaches high levels in the body it causes inflammation and cytotoxicity and it plays an
32
33 important part in fibrotic degeneration, which can contribute to the formation of
34
35 cellulite (7,19-21).
36
37
38
39

40
41 According to Wang (2003), the use of ESWT promotes a cascade of
42
43 physiological events, which have not yet been entirely elucidated. What is known is that
44
45 after the application there is improvement in the metabolism with subsequent stimulus
46
47 of microcirculation by the liberation of nitrous acid and improvement in cell
48
49 permeability, which favors the exchange of substances and the reorganization of the
50
51 cellular matrix. This leads to the rebalance of the production and tamponade of free
52
53 radicals. This data was reaffirmed by the findings of Siems et al. (2005), where there
54
55 was improvement of the aspect of the skin after six sessions of ESWT (9,19,21-23).
56
57
58
59
60

1
2
3 Angehrn et al. (2007) evaluated the effects of ESWT in the treatment of
4 cellulite in 21 women aged between 20 and 60 years. 40 thousand shots were applied in
5 the lateral area of the thigh, twice a week, during six weeks of treatment. The authors
6 used diagnostic ultrasound as evaluation method (Collagenoson®), which evaluated the
7 distribution of the epidermis, dermis and subcutaneous tissue. The results showed that
8 ESWT was effective for the treatment of cellulite, because it promoted the remodeling
9 of the collagen by realigning the dermal fibers and the improvement in the aspect of the
10 skin, which remained latent after six months (8).

11
12 Corroborating the findings of Angehrn et al. (2007), in 2008 two studies
13 evaluated the effects of ESWT in cellulite and body contour. First Kuhn et al. (2008)
14 presented a case study with a 60-year-old woman with cellulite degree three in the
15 gluteus and thighs, where four sessions of ESWT were applied only in the left thigh;
16 after the end of the application the patient underwent orthopedic surgery in the region of
17 the hip in which a sample of tissue of the treated and of the non-treated regions were
18 removed. The authors concluded that there was histological improvement of the dermis
19 with increase of the thickness of the extracellular matrix of the dermis, by means of
20 induction of neocollagen and neo-elastin (8,24).

21
22 The second study was developed by Christ et al. (2008), who evaluated 59
23 women with cellulite, subdivided in two groups in which: the first received six sessions
24 of ESWT for three weeks and the second eight sessions in four weeks of treatment.
25 The results presented were considered excellent, because there was an increase in the
26 elasticity and firmness of the skin in both groups after three and six months of therapy
27 of 95 to 105%, respectively (15).

1
2
3 Angehrn et al. (2007); Kuhn et al. and Christ et al. (2008) found in their
4
5 studies similar results which involve induction of formation of collagen, neo-elastin,
6
7 improvement of the aspect and elasticity of skin and remodeling of collagen (8,15,24).
8

9
10 Collagen is a protein synthesized by the fibroblast, fundamental in the
11
12 constitution of the extracellular matrix and of the conjunctive tissue, and its main
13
14 objective is to structure and protect the properties of the skin. In cellulite, there are
15
16 structural alterations in the intracellular and extracellular matrix, degradation of
17
18 collagen and of fibroblasts. One of the objectives to be achieved is to stimulate the
19
20 fibroblasts to induce the synthesis of new collagen fibers (neocollagen) and elastin (neo-
21
22 elastin) with subsequent restructuring of the properties of the tissue, which we call
23
24 collagen remodeling (14-16,25,26). The cells in our body recognize mechanical stimuli,
25
26 due to this the fibroblasts are activated and start a cascade of formation of neocollagen
27
28 and neo-elastin and the restructuring of the tissue. This mechanism is called
29
30 mechanotransduction of signal, when the cell responds to a mechanical stimulus with
31
32 biochemical reactions in cascade and act on the cellular level transmitting the energy to
33
34 the extracellular matrix (27).
35
36

37
38 The first stimulus of ESWT is mechanical, characterized by the peak of
39
40 pressure, short duration, with high density of energy and low frequency; when in
41
42 contact with the treatment area this mechanical energy leads to the activation of
43
44 mechanotransduction of signal and the extracellular matrix mobilizes the cells. These
45
46 cells, respond to the indirect effect of shock wave cavitation, bringing forth alterations
47
48 beneficial for the treatment of cellulite such as the increase of local circulation and
49
50 stimulus to the production of collagen, leading to the restructuring of the dermis and
51
52 epidermis, thus restoring the elasticity of the conjunctive tissue and improving skin
53
54 texture (2,28).
55
56
57
58
59
60

1
2
3 Adatto et al. (2010) and Adatto et al. (2011) evaluated the effects of ESWT
4 in cellulite and the improvement of body contour, and in their last study the objective
5 was to evaluate ESWT in localized fat (29,30).
6
7

8
9 For the treatment of cellulite, 25 women who received 3000 shots in an area
10 of 10x15cm² in the lateral region of the thigh were evaluated and six sessions twice a
11 week were performed. The results demonstrated reduction of the appearance of cellulitis
12 after three months of the last treatment. The authors concluded that the shockwave
13 therapy can stimulate lymphatic drainage and microcirculation. In the study about
14 improvement of the body contour, 14 women were evaluated after eight sessions in the
15 lateral region of the thigh, with the application of 4500 shots. There was temporary
16 improvement in the texture and elasticity of skin, and a reduction? of the adipose tissue
17 (29,30).
18
19

20
21 Besides the mechanical effects, ESWT has the effects of unstable cavitation,
22 which is the formation of gas microbubbles in the biological liquids, which implode
23 causing physiological effects (3-6). According to Steinbach et al. (1993) and Kuhn et al.
24 (2008), the effects caused by cavitation are related with the dose of energy used to
25 achieve the therapeutic objective. When high doses of energy are used, the damages are
26 the increase of cellular diffusion with improvement in the permeability of the cellular
27 membrane, lesions in the endoplasmic reticulum in the cytoskeleton that gives form to
28 the cell and in the cellular junctions. All these factors may lead to apoptosis. There is
29 still the liberation of endothelial growth factor, synthesis of eNOS enzyme, which
30 induces the production of endothelial nitric oxide (NO) (24,31). For smaller doses of
31 energy, the principle of action of ESWT is the stimulus to the self-regeneration of the
32 tissue; therefore, smaller doses are indicated for the treatment of different types of
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 musculoskeletal diseases. However, both doses stimulate the local blood circulation and
4
5 the metabolism (31).
6

7
8 During the bibliographic revision carried out in the present study, out of the
9
10 12 articles evaluated, most of them used high doses between 3 and 4 bar for devices
11
12 with pneumatic shockwave energy generators. Therefore, we can suggest that, by the
13
14 findings in this study, that high energy is efficient in the treatment of localized fat,
15
16 because the effects of ESWT can lead to cellular apoptosis (28,32,33).
17

18
19 Another Ferraro et al. (2012) study, also demonstrated decrease in thickness
20
21 of the layer of localized fat and of the circumference of the treated area and
22
23 improvement in the aspect of the skin after 12 months of the end of treatment. However,
24
25 the study protocol evaluated the combination of ESWT with cryolipolysis sliding mode,
26
27 using the device that presents both these therapies, but it was not possible to evaluate if
28
29 the results could be attributed solely to the use of ESWT, because cryolipolysis also
30
31 presents significant results for the reduction of localized fat. It is worth mentioning that
32
33 the technique of cryolipolysis sliding mode differs from conventional cryolipolysis
34
35 usually mentioned in literature [33].
36
37

38
39 Knobloch et al. (2013) compared two groups, application of ESWT
40
41 (0,25mJ/mm²) in gluteus and thighs, followed by thigh strengthening training and the
42
43 second control group with application of ESWT (0,01mJ/mm²) in gluteus and thigh,
44
45 followed by thigh and gluteus strengthening training. In both groups, six sessions were
46
47 applied for two weeks. The objective was to evaluate if the muscle strength training
48
49 would lead to additional effects to the ESWT therapy in the improvement of cellulite.
50
51 The results obtained demonstrated that the combination of therapies presented superior
52
53 results in relation to the control group, with significant alterations of 24% in the
54
55 improvement of the aspect of cellulite. The authors suggest that this improvement
56
57
58
59
60

1
2
3 occurred through the mechanical energy of ESWT, which acts on the fibrous septum
4
5 and induces their reorganization, making the skin smoother (34).
6

7 Proving the previous results, Russe-Wilflingseder et al. (2013) evaluated, in
8
9 their double blind randomized clinical trial, the effects of ESWT in the treatment of
10
11 cellulite. A placebo group was evaluated, where the transfer of energy was blocked and
12
13 for a second group that effectively received intervention were performed weekly
14
15 sessions for seven weeks. The treated group presented improvement in the aspect of
16
17 skin, in the appearance of cellulitis with decrease of undulations and depth and
18
19 reduction in the circumference of thighs after three months of treatment; however, the
20
21 researchers suggested that these results can be temporary (28).
22
23

24 The findings corroborate the effects of ESWT in the remodeling of collagen,
25
26 formation of neocollagen and neo-elastin, acting in the improvement of tissue firmness
27
28 and preventing that the irregularities present in the subcutaneous tissue project onto the
29
30 dermis, decreasing the aspect of “orange peel” skin (8,15,30,35).
31
32
33

34 In 2014, Schlaudraff and collaborators evaluated the individual clinical
35
36 result of the treatment of cellulitis with ESWT in 14 women who received 7500 shots in
37
38 unilateral thigh and gluteus in two sessions per week in four weeks, totalizing eight
39
40 treatments. The authors concluded that there was improvement and decrease in the
41
42 degrees of cellulite, but that this result cannot be correlated with individual variables
43
44 such as body mass index (BMI), weight, height or age (28).
45
46

47 Nassar et al. (2015) evaluated the efficacy of lipolysis induced by ESWT, in
48
49 15 individuals; eight sessions during four weeks were carried out, and 4000 shots were
50
51 applied. The authors concluded that ESWT was efficient in the improvement of body
52
53 contour, with reduction of the circumference and the fat layer and improvement of the
54
55 appearance of cellulite after three months of the end of the treatment (36).
56
57
58
59
60

1
2
3 The most recent study published, Hexsel et al. (2016), is in accordance with
4 the previous findings. This study evaluated the efficiency of ESWT in the treatment of
5 cellulite and body contour with 5000 shots in the region of gluteus and posterior region
6 of thigh for 12 sessions. The authors concluded that there was decrease in the
7 circumference of the treated area, measured in the magnetic resonance exam, with
8 improvement of the severity of the degrees of cellulite (37).
9
10
11
12
13
14
15

16 The systematic review carried out in the present study shows that the use of
17 ESWT in the treatment of cellulite and localized fat seems to be a tool with efficient
18 results, provided that the parameters used are observed. The main effects reported in
19 these studies are related with the activation of the mechanisms of action and the
20 physiological effects as a result of the stimulus by mechanical shockwaves and effects
21 of cavitation promoted by the shockwave therapy.
22
23
24
25
26
27
28

29 The main effects observed in the biological tissue are: damage to the
30 extracellular matrix which promotes a cascade of physiological reactions that favor the
31 reorganization of the extracellular medium, increase of blood and lymphatic circulation,
32 alterations in the permeability of the cellular membrane, liberation of nitric oxide,
33 balance of free radicals, drainage of molecular proteins, stimulus for fibroblast
34 activation by mechanotransduction, formation of neocollagen, neo-elastin and
35 remodeling of collagen. Other results of the action of ESWT in the biological tissues
36 were described by Kuhn et al., (2008) and Nassar et al., (2015), the authors suggest that
37 ESWT can induce lipolysis and/or apoptosis of the adipose cell (8,24,36).
38
39
40
41
42
43
44
45
46
47
48

49 The few studies about ESWT in aesthetic affections show that the
50 mechanisms of action and physiological effects of this technique are beneficial and
51 promising for the treatment of cellulite, body contour and localized fat, particularly
52 because they represent a safe and non-invasive therapeutic mode. There are no reports
53
54
55
56
57
58
59
60

1
2
3 of adverse reactions of ESWT in the treatment; only the discomfort reported by the
4 therapist due to the noise emitted by the shock of the projectile colliding with the tip of
5 the applicator; to minimize this discomfort, the use of an auricular protector is
6 suggested.
7
8
9
10

11 12 13 14 **Conclusion**

15
16 The results presented in this study show the efficacy of the ESWT technique
17 as a safe and non-invasive method. Its mode of action in the biological tissue benefits
18 the treatment of cellulite, body contour and localized fat. Because it is a relatively new
19 therapy in the aesthetic area, ESWT is a promising field for studies and application, but
20 it still requires investigation as to the mechanisms of action and their interaction with
21 the adipose tissue, which demands further experimental studies with ESWT in aesthetic
22 affections.
23
24
25
26
27
28
29
30
31
32
33

34 **Conflicts of Interest**

35
36 The authors declare that there are no conflicts of interest regarding the
37 publication of this paper.
38
39
40
41
42

43 **References**

- 44
45
46 1. Quatresooz P, Xhaufaire-Uhoda E, Piérard-Franchimont, Piérard GE. Cellulite
47 histopathology and related mechanobiology. Int. J. Cosmet. Sci 2000; 28 (3): 207–
48 210, 2006.
49
50
51
52
53 2. Thiel M, Nieswand M, Dörffel M. Review: The use of shock waves in medicine a tool
54 of the modern OR: An overview of basic physical principles, history and research.
55 Minim. Invasive Ther. Allied Technol 2000; 9(3):247-253.
56
57
58
59
60

- 1
2
3 3. Notarnicola A, Moretti B. The biological effects of extracorporeal shock wave therapy
4 (eswt) on tendon tissue. *Muscles. Ligaments Tendons J* 2012;2(1):33-37, 2012.
- 5
6
7 4. Ogden JA, Tóth-Kischkat A, Schultheiss R. Principles of shock wave therapy. *Clin.*
8
9 *Orthop. Relat. Rel* 2001; 387:8–17.
- 10
11 5. Kertzman P, Lenza M, Pedrinelli A, Ejnisman B. Tratamento por ondas de choque
12 nas doenças musculoesqueléticas e consolidação óssea – Análise qualitativa da
13 literature. *Rev. Bras. Ortop* 2010;50 (1): 3–8.
- 14
15
16 6. Wang C. Extracorporeal shockwave therapy in musculoskeletal disorders. *J. Orthop.*
17
18 *Surg. Res* 2012;7(1):7-11.
- 19
20
21 7. Knobloch K, Kraemer R. Extracorporeal shock wave therapy (ESWT) for the
22 treatment of cellulite-A current metaanalysis. *Int. J. Surg* 2015; 24:1-8.
- 23
24
25 8. Angehrn F, Kuhn C, Voss A. Can cellulite be treated with low-energy extracorporeal
26 shock wave therapy?. *Clin. Interv. Aging* 2007; 2(4): 623- 630.
- 27
28
29 9. Wang C. An overview of shock wave therapy in musculoskeletal disorders. *Chang*
30
31 *Gung Med. J* 2003; 26(4):220-232.
- 32
33
34 10. Malliaropoulos N, Crate G, Meke M, Korakakis V, Nauck T, Lohrer H, et al.
35
36 Success and Recurrence Rate after Radial Extracorporeal Shock Wave Therapy for
37
38 Plantar Fasciopathy: A Retrospective Study. *Biomed Res Int* 2016; 1-8.
- 39
40
41 11. Chen TW, Lin CW, Lee CL, Chen CH, Chen YJ, Lin TY, et al. The efficacy of
42 shock wave therapy in patients with knee osteoarthritis and popliteal cyamella.
43
44 *Kaohsiung J. Med. Sci* 2014;30(7): 362-370.
- 45
46
47 12. Rossi AB, Vergnanini AL. Cellulite: a review. *J. Eur. Acad. Dermatol. Venereol*
48
49
50 2000;14(4):251-262.
- 51
52
53 13. Avram MM. Cellulite: a review of its physiology and treatment. *J. Cosmet. Laser*
54
55
56 *Ther* 2004; 6(4):181-185.
- 57
58
59
60

14. Afonso MJPJ, Tucunduva MCT, Pinheiro BVM, Bagatin E. Cellulite: a review. *Surg Cosmet Dermatol* 2010;2(3): 214-219.
15. Christ C, Brenke R, Sattler G, Siems W, Novak P, Daser A. Improvement in Skin Elasticity in the Treatment of Cellulite and Connective Tissue Weakness by Means of Extracorporeal Pulse Activation Therapy. *Aesthetic Surg. J* 2008; 28(5): 538- 544.
16. Rawlings AV. Cellulite and its treatment. *International Journal of Cosmetic Science* 2006;28(3):175-190.
17. Writers AM. Cellulite: No clear evidence that any type of treatment is effective. *Drugs Ther Perspect* 2015;31(12): 437-440.
18. Zerini I, Sisti A, Cuomo R, Ciappi S, Russo F, Brandi C, et al. Cellulite treatment: a comprehensive literature review. *Journal of Cosmetic Dermatology* 2015;14(3):224-240.
19. Siems W, Grune T, Voss P, Brenke R. Anti-fibrosclerotic effects of shock wave therapy in lipedema and cellulite. *Biofactors* 2005; 24(1): 275-282.
20. Kandola K, Bowman A. Oxidative stress – a key emerging impact factor in health, ageing, lifestyle and aesthetics. *International Journal of Cosmetic Science* 2015; 37:1-8.
21. Koshiyama K, Kodama T, Yano T, Fujikawa S. Structural Change in Lipid Bilayers and Water Penetration Induced by Shock Waves: Molecular Dynamics Simulations. *Biophysical Journal* 2006; 91(6): 2198- 2205.
22. Moosavi-nejad SJ, Satoh M, Takayama K, Island G. Shock wave induced cytoskeletal and morphological deformations in a human renal carcinoma cell line. *Cancer Science* 2006; 97(4): 296-304.
23. Frairia R, Berta L. Biological Effects of Extracorporeal Shock Waves on Fibroblasts. A Review. *Muscles Ligaments Tendons* 2011; 1(4): 138-147.

- 1
2
3 24. Kuhn C. Impact of extracorporeal shock waves on the human skin with cellulite: A
4 case study of an unique instance. *Clinical Interventions in Aging* 2008; 3(1): 201-210.
5
6
7 25. Baxter SC, Morales EMO, Goldsmith EC. Adaptive Changes in Cardiac Fibroblast
8 Morphology and Collagen Organization as a Result of Mechanical Environment. *Cell*
9 *Biochemistry and Biophysics* 2008; 51(1): 33–44.
10
11
12
13 26. Khan MH, Victor F, Rao B, Sadick NS. Treatment of cellulite. *J. Am. Dermatology*
14 2010; 62(3): 361-370.
15
16
17
18 27. Humphrey JD, Dufresne ER, Schwartz MA. Mechanotransduction and extracellular
19 matrix homeostasis. *Nat Rev Mol Cell Biol* 2014;15(12): 802-812.
20
21
22
23 28. Schlaudraff KU, Kiessling MC, Császár NBM, Schmitz C. Predictability of the
24 individual clinical outcome of extracorporeal shock wave therapy for cellulite. *Clin.*
25 *Cosmet. Investig. Dermatol* 2014; 7: 171-182.
26
27
28
29 29. Adatto M, Adatto NR, Servant J, Vester J, Novak P, Krotz A. Controlled,
30 randomized study evaluating the effects of treating cellulite with AWT® / EPAT®.
31 *Journal of Cosmetic and Laser Therapy* 2010; 12: 176-182.
32
33
34
35 30. Adatto MA, Adatto NR, Novak P, Krotz A, Haller G. Body shaping with acoustic
36 wave therapy AWT (®) /EPAT (®): randomized, controlled study on 14 subjects. *J.*
37 *Cosmet. Laser Ther* 2011; 13: 291-296.
38
39
40
41 31. Steinbach P, Hofstaedter F, Nicolai H, Roessler W, Wieland W. Determination of the
42 energy-dependent extent of vascular damage caused by high-energy shock waves in an
43 umbilical cord model. *Urological Research* 1993; 21: 279-282.
44
45
46
47
48 32. Butterworth PA, Walsh TP, Pennisi YD, Chesne AD, Schmitz C, Nancarrow SA.
49 The effectiveness of extracorporeal shock wave therapy for the treatment of lower
50 limb ulceration: a systematic review. *Journal of Foot and Ankle Research* 2015; 8:1-9.
51
52
53
54
55
56
57
58
59
60

- 1
2
3 33. Ferraro GA, De Francesco F, Cataldo C, Rossano F, Nicoletti G, D'Andrea.
4 Synergistic effects of cryolipolysis and shock waves for noninvasive body contouring.
5 Aesthetic Plast. Surg 2012; 36 (3):666-679.
6
7
8
9
10 34. Knobloch K, Joest B, Krämer R, Vogt PM. Cellulite and focused extracorporeal
11 shockwave therapy for non-invasive body contouring: A randomized trial. Dermatol.
12 Ther 2013;3(2): 143-155.
13
14
15
16 35. Nishida C. Appropriate body-mass index for Asian populations and its implications
17 for policy and intervention strategies. Lancet 2004; 363(9403): 157-163.
18
19
20
21 36. Nassar AH, Dorizas AS, Shafai A, Sadick NS. A randomized, controlled clinical
22 study to investigate the safety and efficacy of acoustic wave therapy in body
23 contouring. Dermatol. Surg 2015; 41: 366-370.
24
25
26
27 37. Hexsel D, Camozzato FO, Silva AF, Siega C, Camozzato FO. Acoustic wave
28 therapy for cellulite, body shaping and fat reduction and fat reduction. Journal of
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only




Figure 1. Electromagnetic System (a): (A) coils, (B) projectile, (C) tip of the applicator. Pneumatic System (b): (B) projectile; (C) tip of the applicator, (D) compressed air inlet.

Table 1. Published studies about the action of ESWT on cellulite, body contour and localized fat.

Author	Year	Protocol	Treatment Area	Conclusion
Siems et al.	2005	ESWT + Decongestive therapy	Thigh	Decrease of oxidative stress and improvement of skin properties.
Angehrt et al.	2007	ESWT	Lateral region of thigh	Remodeling of collagen and improvement of the aspect of skin.
Kuhn et al.	2008	ESWT	Thigh and gluteus	Histological improvement of epidermis and of the extracellular matrix of the dermis.
Christ et al.	2008	ESWT	Thigh and gluteus	Improvement of elasticity and tightness of skin.
Adatto et al.	2010 2011	ESWT	Lateral region of the thigh	Improvement in skin texture and elasticity, number of depression and decrease of thickness of the subcutaneous tissue.
Ferraro et al.	2012	ESWT+ Cryolipolysis	Abdomen, arms, gluteus and thigh.	Decrease of the thickness of the adipose tissue, circumference of the treated area and improvement of the aspect of the skin.
Knoblock et al.	2013	ESWT+ Muscle strengthening.	Thigh and gluteus	Improvement of the aspect of cellulite.
Russelwifflingseder et al.	2013	ESWT	Thigh	Temporary improvement of the aspect of the skin and reduction in the circumference of the thigh.
Schlaudraff et al.	2014	ESWT	Thigh and gluteus	Decrease in the degree of cellulitis.
Nassar et al.	2015	ESWT	Thigh	Reduction of the circumference of the treated area, thickness of the fat layer and appearance of cellulite.
Hexsel et al.	2016	ESWT	Thigh and gluteus	Decrease of the circumference of the treated area and severity of the degrees of cellulite.